

# The Hahn-Banach Theorem for Real Vector Spaces

Gertrud Bauer  
<http://www.in.tum.de/~bauerg/>

October 10, 2011

## Abstract

The Hahn-Banach Theorem is one of the most fundamental results in functional analysis. We present a fully formal proof of two versions of the theorem, one for general linear spaces and another for normed spaces. This development is based on simply-typed classical set-theory, as provided by Isabelle/HOL.

## Contents


<b>1</b>	<b>Preface</b>	<b>3</b>
<b>I</b>	<b>Basic Notions</b>	<b>4</b>
<b>2</b>	<b>Bounds</b>	<b>4</b>
<b>3</b>	<b>Vector spaces</b>	<b>4</b>
3.1	Signature . . . . .	5
3.2	Vector space laws . . . . .	5
<b>4</b>	<b>Subspaces</b>	<b>8</b>
4.1	Definition . . . . .	8
4.2	Linear closure . . . . .	9
4.3	Sum of two vectorspaces . . . . .	10
4.4	Direct sums . . . . .	11
<b>5</b>	<b>Normed vector spaces</b>	<b>12</b>
5.1	Quasinorms . . . . .	12
5.2	Norms . . . . .	12
5.3	Normed vector spaces . . . . .	13
<b>6</b>	<b>Linearforms</b>	<b>13</b>

<b>7</b>	<b>An order on functions</b>	<b>14</b>
7.1	The graph of a function . . . . .	14
7.2	Functions ordered by domain extension . . . . .	14
7.3	Domain and function of a graph . . . . .	15
7.4	Norm-preserving extensions of a function . . . . .	15
<b>8</b>	<b>The norm of a function</b>	<b>16</b>
8.1	Continuous linear forms . . . . .	16
8.2	The norm of a linear form . . . . .	16
<b>9</b>	<b>Zorn's Lemma</b>	<b>18</b>
<b>II</b>	<b>Lemmas for the Proof</b>	<b>19</b>
<b>10</b>	<b>The supremum w.r.t. the function order</b>	<b>19</b>
<b>11</b>	<b>Extending non-maximal functions</b>	<b>21</b>
<b>III</b>	<b>The Main Proof</b>	<b>23</b>
<b>12</b>	<b>The Hahn-Banach Theorem</b>	<b>23</b>
12.1	The Hahn-Banach Theorem for vector spaces . . . . .	23
12.2	Alternative formulation . . . . .	23
12.3	The Hahn-Banach Theorem for normed spaces . . . . .	24

# 1 Preface

This is a fully formal proof of the Hahn-Banach Theorem. It closely follows the informal presentation given in Heuser's textbook [1, § 36]. Another formal proof of the same theorem has been done in Mizar [3]. A general overview of the relevance and history of the Hahn-Banach Theorem is given by Narici and Beckenstein [2].

The document is structured as follows. The first part contains definitions of basic notions of linear algebra: vector spaces, subspaces, normed spaces, continuous linear-forms, norm of functions and an order on functions by domain extension. The second part contains some lemmas about the supremum (w.r.t. the function order) and extension of non-maximal functions. With these preliminaries, the main proof of the theorem (in its two versions) is conducted in the third part. The dependencies of individual theories are as follows.



## Part I

# Basic Notions

## 2 Bounds

```

theory Bounds
imports Main ~~/src/HOL/Library/ContNotDenum
begin

locale lub =
  fixes A and x
  assumes least [intro?]: ( $\bigwedge a. a \in A \implies a \leq b$ )  $\implies x \leq b$ 
  and upper [intro?]:  $a \in A \implies a \leq x$ 

lemmas [elim?] = lub.least lub.upper

definition the-lub :: 'a::order set  $\Rightarrow$  'a
  where the-lub A = The (lub A)

notation (xsymbols)
  the-lub ( $\bigsqcup$  - [90] 90)

lemma the-lub-equality [elim?]:
  assumes lub A x
  shows  $\bigsqcup A = (x::'a::order)$ 
  <proof>

lemma the-lubI-ex:
  assumes ex:  $\exists x. \text{lub } A \ x$ 
  shows lub A ( $\bigsqcup A$ )
  <proof>

lemma lub-compat: lub A x = isLub UNIV A x
  <proof>

lemma real-complete:
  fixes A :: real set
  assumes nonempty:  $\exists a. a \in A$ 
  and ex-upper:  $\exists y. \forall a \in A. a \leq y$ 
  shows  $\exists x. \text{lub } A \ x$ 
  <proof>

end

```

## 3 Vector spaces

```

theory Vector-Space
imports Complex-Main Bounds ~~/src/HOL/Library/Zorn
begin

```

### 3.1 Signature

For the definition of real vector spaces a type  $'a$  of the sort  $\{plus, minus, zero\}$  is considered, on which a real scalar multiplication  $\cdot$  is declared.

**consts**

$prod :: real \Rightarrow 'a::\{plus, minus, zero\} \Rightarrow 'a$  (infixr  $'(*)$  70)

**notation** (*xsymbols*)

$prod$  (infixr  $\cdot$  70)

**notation** (*HTML output*)

$prod$  (infixr  $\cdot$  70)

### 3.2 Vector space laws

A *vector space* is a non-empty set  $V$  of elements from  $'a$  with the following vector space laws: The set  $V$  is closed under addition and scalar multiplication, addition is associative and commutative;  $-x$  is the inverse of  $x$  w. r. t. addition and  $0$  is the neutral element of addition. Addition and multiplication are distributive; scalar multiplication is associative and the real number  $1$  is the neutral element of scalar multiplication.

**locale**  $var-V = \text{fixes } V$

**locale**  $vectorspace = var-V +$

**assumes** *non-empty* [*iff, intro?*]:  $V \neq \{\}$

**and** *add-closed* [*iff*]:  $x \in V \Longrightarrow y \in V \Longrightarrow x + y \in V$

**and** *mult-closed* [*iff*]:  $x \in V \Longrightarrow a \cdot x \in V$

**and** *add-assoc*:  $x \in V \Longrightarrow y \in V \Longrightarrow z \in V \Longrightarrow (x + y) + z = x + (y + z)$

**and** *add-commute*:  $x \in V \Longrightarrow y \in V \Longrightarrow x + y = y + x$

**and** *diff-self* [*simp*]:  $x \in V \Longrightarrow x - x = 0$

**and** *add-zero-left* [*simp*]:  $x \in V \Longrightarrow 0 + x = x$

**and** *add-mult-distrib1*:  $x \in V \Longrightarrow y \in V \Longrightarrow a \cdot (x + y) = a \cdot x + a \cdot y$

**and** *add-mult-distrib2*:  $x \in V \Longrightarrow (a + b) \cdot x = a \cdot x + b \cdot x$

**and** *mult-assoc*:  $x \in V \Longrightarrow (a * b) \cdot x = a \cdot (b \cdot x)$

**and** *mult-1* [*simp*]:  $x \in V \Longrightarrow 1 \cdot x = x$

**and** *negate-eq1*:  $x \in V \Longrightarrow -x = (-1) \cdot x$

**and** *diff-eq1*:  $x \in V \Longrightarrow y \in V \Longrightarrow x - y = x + -y$

**begin**

**lemma** *negate-eq2*:  $x \in V \Longrightarrow (-1) \cdot x = -x$

*<proof>*

**lemma** *negate-eq2a*:  $x \in V \Longrightarrow -1 \cdot x = -x$

*<proof>*

**lemma** *diff-eq2*:  $x \in V \Longrightarrow y \in V \Longrightarrow x + -y = x - y$

*<proof>*

**lemma** *diff-closed* [*iff*]:  $x \in V \Longrightarrow y \in V \Longrightarrow x - y \in V$

*<proof>*

**lemma** *neg-closed* [*iff*]:  $x \in V \Longrightarrow -x \in V$

*<proof>*

**lemma** *add-left-commute*:  $x \in V \implies y \in V \implies z \in V \implies x + (y + z) = y + (x + z)$   
 ⟨*proof*⟩

**theorems** *add-ac = add-assoc add-commute add-left-commute*

The existence of the zero element of a vector space follows from the non-emptiness of carrier set.

**lemma** *zero [iff]*:  $0 \in V$   
 ⟨*proof*⟩

**lemma** *add-zero-right [simp]*:  $x \in V \implies x + 0 = x$   
 ⟨*proof*⟩

**lemma** *mult-assoc2*:  $x \in V \implies a \cdot b \cdot x = (a * b) \cdot x$   
 ⟨*proof*⟩

**lemma** *diff-mult-distrib1*:  $x \in V \implies y \in V \implies a \cdot (x - y) = a \cdot x - a \cdot y$   
 ⟨*proof*⟩

**lemma** *diff-mult-distrib2*:  $x \in V \implies (a - b) \cdot x = a \cdot x - (b \cdot x)$   
 ⟨*proof*⟩

**lemmas** *distrib =*  
*add-mult-distrib1 add-mult-distrib2*  
*diff-mult-distrib1 diff-mult-distrib2*

Further derived laws:

**lemma** *mult-zero-left [simp]*:  $x \in V \implies 0 \cdot x = 0$   
 ⟨*proof*⟩

**lemma** *mult-zero-right [simp]*:  $a \cdot 0 = (0::'a)$   
 ⟨*proof*⟩

**lemma** *minus-mult-cancel [simp]*:  $x \in V \implies (- a) \cdot - x = a \cdot x$   
 ⟨*proof*⟩

**lemma** *add-minus-left-eq-diff*:  $x \in V \implies y \in V \implies - x + y = y - x$   
 ⟨*proof*⟩

**lemma** *add-minus* [simp]:  $x \in V \implies x + - x = 0$   
 ⟨*proof*⟩

**lemma** *add-minus-left* [simp]:  $x \in V \implies - x + x = 0$   
 ⟨*proof*⟩

**lemma** *minus-minus* [simp]:  $x \in V \implies - (- x) = x$   
 ⟨*proof*⟩

**lemma** *minus-zero* [simp]:  $- (0::'a) = 0$   
 ⟨*proof*⟩

**lemma** *minus-zero-iff* [simp]:

**assumes**  $x: x \in V$   
**shows**  $(-x = 0) = (x = 0)$   
 ⟨proof⟩

**lemma** *add-minus-cancel* [simp]:  $x \in V \implies y \in V \implies x + (-x + y) = y$   
 ⟨proof⟩

**lemma** *minus-add-cancel* [simp]:  $x \in V \implies y \in V \implies -x + (x + y) = y$   
 ⟨proof⟩

**lemma** *minus-add-distrib* [simp]:  $x \in V \implies y \in V \implies -(x + y) = -x + -y$   
 ⟨proof⟩

**lemma** *diff-zero* [simp]:  $x \in V \implies x - 0 = x$   
 ⟨proof⟩

**lemma** *diff-zero-right* [simp]:  $x \in V \implies 0 - x = -x$   
 ⟨proof⟩

**lemma** *add-left-cancel*:  
**assumes**  $x: x \in V$  **and**  $y: y \in V$  **and**  $z: z \in V$   
**shows**  $(x + y = x + z) = (y = z)$   
 ⟨proof⟩

**lemma** *add-right-cancel*:  $x \in V \implies y \in V \implies z \in V \implies (y + x = z + x) = (y = z)$   
 ⟨proof⟩

**lemma** *add-assoc-cong*:  
 $x \in V \implies y \in V \implies x' \in V \implies y' \in V \implies z \in V$   
 $\implies x + y = x' + y' \implies x + (y + z) = x' + (y' + z)$   
 ⟨proof⟩

**lemma** *mult-left-commute*:  $x \in V \implies a \cdot b \cdot x = b \cdot a \cdot x$   
 ⟨proof⟩

**lemma** *mult-zero-uniq*:  
**assumes**  $x: x \in V$   $x \neq 0$  **and**  $ax: a \cdot x = 0$   
**shows**  $a = 0$   
 ⟨proof⟩

**lemma** *mult-left-cancel*:  
**assumes**  $x: x \in V$  **and**  $y: y \in V$  **and**  $a: a \neq 0$   
**shows**  $(a \cdot x = a \cdot y) = (x = y)$   
 ⟨proof⟩

**lemma** *mult-right-cancel*:  
**assumes**  $x: x \in V$  **and**  $neg: x \neq 0$   
**shows**  $(a \cdot x = b \cdot x) = (a = b)$   
 ⟨proof⟩

**lemma** *eq-diff-eq*:  
**assumes**  $x: x \in V$  **and**  $y: y \in V$  **and**  $z: z \in V$   
**shows**  $(x = z - y) = (x + y = z)$

*<proof>*

**lemma** *add-minus-eq-minus*:

**assumes**  $x: x \in V$  **and**  $y: y \in V$  **and**  $xy: x + y = 0$

**shows**  $x = -y$

*<proof>*

**lemma** *add-minus-eq*:

**assumes**  $x: x \in V$  **and**  $y: y \in V$  **and**  $xy: x - y = 0$

**shows**  $x = y$

*<proof>*

**lemma** *add-diff-swap*:

**assumes**  $vs: a \in V$   $b \in V$   $c \in V$   $d \in V$

**and**  $eq: a + b = c + d$

**shows**  $a - c = d - b$

*<proof>*

**lemma** *vs-add-cancel-21*:

**assumes**  $vs: x \in V$   $y \in V$   $z \in V$   $u \in V$

**shows**  $(x + (y + z) = y + u) = (x + z = u)$

*<proof>*

**lemma** *add-cancel-end*:

**assumes**  $vs: x \in V$   $y \in V$   $z \in V$

**shows**  $(x + (y + z) = y) = (x = -z)$

*<proof>*

**end**

**end**

## 4 Subspaces

**theory** *Subspace*

**imports** *Vector-Space*  $\sim\sim$  */src/HOL/Library/Set-Algebras*

**begin**

### 4.1 Definition

A non-empty subset  $U$  of a vector space  $V$  is a *subspace* of  $V$ , iff  $U$  is closed under addition and scalar multiplication.

**locale** *subspace* =

**fixes**  $U :: 'a::\{minus, plus, zero, uminus\}$  *set* **and**  $V$

**assumes** *non-empty* [*iff, intro*]:  $U \neq \{\}$

**and** *subset* [*iff*]:  $U \subseteq V$

**and** *add-closed* [*iff*]:  $x \in U \implies y \in U \implies x + y \in U$

**and** *mult-closed* [*iff*]:  $x \in U \implies a \cdot x \in U$

**notation** (*symbols*)

*subspace* (**infix**  $\leq 50$ )

**declare** *vectorspace.intro* [*intro?*] *subspace.intro* [*intro?*]

**lemma** *subspace-subset* [*elim*]:  $U \trianglelefteq V \implies U \subseteq V$   
 ⟨*proof*⟩

**lemma** (**in** *subspace*) *subsetD* [*iff*]:  $x \in U \implies x \in V$   
 ⟨*proof*⟩

**lemma** *subspaceD* [*elim*]:  $U \trianglelefteq V \implies x \in U \implies x \in V$   
 ⟨*proof*⟩

**lemma** *rev-subspaceD* [*elim?*]:  $x \in U \implies U \trianglelefteq V \implies x \in V$   
 ⟨*proof*⟩

**lemma** (**in** *subspace*) *diff-closed* [*iff*]:  
**assumes** *vectorspace*  $V$   
**assumes**  $x: x \in U$  **and**  $y: y \in U$   
**shows**  $x - y \in U$   
 ⟨*proof*⟩

Similar as for linear spaces, the existence of the zero element in every subspace follows from the non-emptiness of the carrier set and by vector space laws.

**lemma** (**in** *subspace*) *zero* [*intro*]:  
**assumes** *vectorspace*  $V$   
**shows**  $0 \in U$   
 ⟨*proof*⟩

**lemma** (**in** *subspace*) *neg-closed* [*iff*]:  
**assumes** *vectorspace*  $V$   
**assumes**  $x: x \in U$   
**shows**  $-x \in U$   
 ⟨*proof*⟩

Further derived laws: every subspace is a vector space.

**lemma** (**in** *subspace*) *vectorspace* [*iff*]:  
**assumes** *vectorspace*  $V$   
**shows** *vectorspace*  $U$   
 ⟨*proof*⟩

The subspace relation is reflexive.

**lemma** (**in** *vectorspace*) *subspace-refl* [*intro*]:  $V \trianglelefteq V$   
 ⟨*proof*⟩

The subspace relation is transitive.

**lemma** (**in** *vectorspace*) *subspace-trans* [*trans*]:  
 $U \trianglelefteq V \implies V \trianglelefteq W \implies U \trianglelefteq W$   
 ⟨*proof*⟩

## 4.2 Linear closure

The *linear closure* of a vector  $x$  is the set of all scalar multiples of  $x$ .

**definition**  $lin :: ('a::\{minus, plus, zero\}) \Rightarrow 'a \text{ set}$   
**where**  $lin\ x = \{a \cdot x \mid a. True\}$

**lemma**  $linI$  [intro]:  $y = a \cdot x \Longrightarrow y \in lin\ x$   
 ⟨proof⟩

**lemma**  $linI'$  [iff]:  $a \cdot x \in lin\ x$   
 ⟨proof⟩

**lemma**  $linE$  [elim]:  $x \in lin\ v \Longrightarrow (\bigwedge a::real. x = a \cdot v \Longrightarrow C) \Longrightarrow C$   
 ⟨proof⟩

Every vector is contained in its linear closure.

**lemma** (in *vectorspace*)  $x\text{-lin-}x$  [iff]:  $x \in V \Longrightarrow x \in lin\ x$   
 ⟨proof⟩

**lemma** (in *vectorspace*)  $0\text{-lin-}x$  [iff]:  $x \in V \Longrightarrow 0 \in lin\ x$   
 ⟨proof⟩

Any linear closure is a subspace.

**lemma** (in *vectorspace*)  $lin\text{-subspace}$  [intro]:  
**assumes**  $x: x \in V$   
**shows**  $lin\ x \leq V$   
 ⟨proof⟩

Any linear closure is a vector space.

**lemma** (in *vectorspace*)  $lin\text{-vectorspace}$  [intro]:  
**assumes**  $x \in V$   
**shows** *vectorspace* ( $lin\ x$ )  
 ⟨proof⟩

### 4.3 Sum of two vectorspaces

The *sum* of two vectorspaces  $U$  and  $V$  is the set of all sums of elements from  $U$  and  $V$ .

**lemma**  $sum\text{-def}$ :  $U \oplus V = \{u + v \mid u \in U \wedge v \in V\}$   
 ⟨proof⟩

**lemma**  $sumE$  [elim]:  
 $x \in U \oplus V \Longrightarrow (\bigwedge u \in U, v \in V. x = u + v \Longrightarrow C) \Longrightarrow C$   
 ⟨proof⟩

**lemma**  $sumI$  [intro]:  
 $u \in U \Longrightarrow v \in V \Longrightarrow x = u + v \Longrightarrow x \in U \oplus V$   
 ⟨proof⟩

**lemma**  $sumI'$  [intro]:  
 $u \in U \Longrightarrow v \in V \Longrightarrow u + v \in U \oplus V$   
 ⟨proof⟩

$U$  is a subspace of  $U \oplus V$ .

**lemma**  $subspace\text{-sumI}$  [iff]:

**assumes** *vectorspace*  $U$  *vectorspace*  $V$   
**shows**  $U \leq U \oplus V$   
 ⟨*proof*⟩

The sum of two subspaces is again a subspace.

**lemma** *sum-subspace* [*intro?*]:  
**assumes** *subspace*  $U$   $E$  *vectorspace*  $E$  *subspace*  $V$   $E$   
**shows**  $U \oplus V \leq E$   
 ⟨*proof*⟩

The sum of two subspaces is a vectorspace.

**lemma** *sum-vs* [*intro?*]:  
 $U \leq E \implies V \leq E \implies \text{vectorspace } E \implies \text{vectorspace } (U \oplus V)$   
 ⟨*proof*⟩

## 4.4 Direct sums

The sum of  $U$  and  $V$  is called *direct*, iff the zero element is the only common element of  $U$  and  $V$ . For every element  $x$  of the direct sum of  $U$  and  $V$  the decomposition in  $x = u + v$  with  $u \in U$  and  $v \in V$  is unique.

**lemma** *decomp*:  
**assumes** *vectorspace*  $E$  *subspace*  $U$   $E$  *subspace*  $V$   $E$   
**assumes** *direct*:  $U \cap V = \{0\}$   
**and**  $u1: u1 \in U$  **and**  $u2: u2 \in U$   
**and**  $v1: v1 \in V$  **and**  $v2: v2 \in V$   
**and** *sum*:  $u1 + v1 = u2 + v2$   
**shows**  $u1 = u2 \wedge v1 = v2$   
 ⟨*proof*⟩

An application of the previous lemma will be used in the proof of the Hahn-Banach Theorem (see page ??): for any element  $y + a \cdot x_0$  of the direct sum of a vectorspace  $H$  and the linear closure of  $x_0$  the components  $y \in H$  and  $a$  are uniquely determined.

**lemma** *decomp-H'*:  
**assumes** *vectorspace*  $E$  *subspace*  $H$   $E$   
**assumes**  $y1: y1 \in H$  **and**  $y2: y2 \in H$   
**and**  $x': x' \notin H$   $x' \in E$   $x' \neq 0$   
**and** *eq*:  $y1 + a1 \cdot x' = y2 + a2 \cdot x'$   
**shows**  $y1 = y2 \wedge a1 = a2$   
 ⟨*proof*⟩

Since for any element  $y + a \cdot x'$  of the direct sum of a vectorspace  $H$  and the linear closure of  $x'$  the components  $y \in H$  and  $a$  are unique, it follows from  $y \in H$  that  $a = 0$ .

**lemma** *decomp-H'-H*:  
**assumes** *vectorspace*  $E$  *subspace*  $H$   $E$   
**assumes**  $t: t \in H$   
**and**  $x': x' \notin H$   $x' \in E$   $x' \neq 0$   
**shows** (*SOME*  $(y, a). t = y + a \cdot x' \wedge y \in H$ ) =  $(t, 0)$   
 ⟨*proof*⟩

The components  $y \in H$  and  $a$  in  $y + a \cdot x'$  are unique, so the function  $h'$  defined by  $h'(y + a \cdot x') = h y + a \cdot \xi$  is definite.

```

lemma h'-definite:
  fixes H
  assumes h'-def:
     $h' \equiv \lambda x.$ 
       $let (y, a) = SOME (y, a). (x = y + a \cdot x' \wedge y \in H)$ 
       $in (h\ y) + a * xi$ 
    and  $x: x = y + a \cdot x'$ 
  assumes vectorspace E subspace H E
  assumes  $y: y \in H$ 
  and  $x': x' \notin H\ x' \in E\ x' \neq 0$ 
  shows  $h' x = h\ y + a * xi$ 
   $\langle proof \rangle$ 

end

```

## 5 Normed vector spaces

```

theory Normed-Space
imports Subspace
begin

```

### 5.1 Quasinorms

A *seminorm*  $\|\cdot\|$  is a function on a real vector space into the reals that has the following properties: it is positive definite, absolute homogenous and subadditive.

```

locale norm-syntax =
  fixes  $norm :: 'a \Rightarrow real$  ( $\|\cdot\|$ )

locale seminorm = var-V + norm-syntax +
  constrains  $V :: 'a::\{minus, plus, zero, uminus\}$  set
  assumes ge-zero [iff?]:  $x \in V \Longrightarrow 0 \leq \|x\|$ 
  and abs-homogenous [iff?]:  $x \in V \Longrightarrow \|a \cdot x\| = |a| * \|x\|$ 
  and subadditive [iff?]:  $x \in V \Longrightarrow y \in V \Longrightarrow \|x + y\| \leq \|x\| + \|y\|$ 

declare seminorm.intro [intro?]

```

```

lemma (in seminorm) diff-subadditive:
  assumes vectorspace V
  shows  $x \in V \Longrightarrow y \in V \Longrightarrow \|x - y\| \leq \|x\| + \|y\|$ 
   $\langle proof \rangle$ 

```

```

lemma (in seminorm) minus:
  assumes vectorspace V
  shows  $x \in V \Longrightarrow \|- x\| = \|x\|$ 
   $\langle proof \rangle$ 

```

### 5.2 Norms

A *norm*  $\|\cdot\|$  is a seminorm that maps only the  $0$  vector to  $0$ .

```

locale norm = seminorm +

```

**assumes** *zero-iff* [*iff*]:  $x \in V \implies (\|x\| = 0) = (x = 0)$

### 5.3 Normed vector spaces

A vector space together with a norm is called a *normed space*.

**locale** *normed-vectorspace* = *vectorspace* + *norm*

**declare** *normed-vectorspace.intro* [*intro?*]

**lemma** (**in** *normed-vectorspace*) *gt-zero* [*intro?*]:

**assumes**  $x: x \in V$  **and** *neq*:  $x \neq 0$

**shows**  $0 < \|x\|$

*<proof>*

Any subspace of a normed vector space is again a normed vectorspace.

**lemma** *subspace-normed-vs* [*intro?*]:

**fixes**  $F E$  *norm*

**assumes** *subspace*  $F E$  *normed-vectorspace*  $E$  *norm*

**shows** *normed-vectorspace*  $F$  *norm*

*<proof>*

**end**

## 6 Linearforms

**theory** *Linearform*

**imports** *Vector-Space*

**begin**

A *linear form* is a function on a vector space into the reals that is additive and multiplicative.

**locale** *linearform* =

**fixes**  $V :: 'a::\{minus, plus, zero, uminus\}$  *set* **and**  $f$

**assumes** *add* [*iff*]:  $x \in V \implies y \in V \implies f (x + y) = f x + f y$

**and** *mult* [*iff*]:  $x \in V \implies f (a \cdot x) = a * f x$

**declare** *linearform.intro* [*intro?*]

**lemma** (**in** *linearform*) *neg* [*iff*]:

**assumes** *vectorspace*  $V$

**shows**  $x \in V \implies f (- x) = - f x$

*<proof>*

**lemma** (**in** *linearform*) *diff* [*iff*]:

**assumes** *vectorspace*  $V$

**shows**  $x \in V \implies y \in V \implies f (x - y) = f x - f y$

*<proof>*

Every linear form yields 0 for the 0 vector.

**lemma** (**in** *linearform*) *zero* [*iff*]:

**assumes** *vectorspace*  $V$

```

  shows  $f 0 = 0$ 
  <proof>

end

```

## 7 An order on functions

```

theory Function-Order
imports Subspace Linearform
begin

```

### 7.1 The graph of a function

We define the *graph* of a (real) function  $f$  with domain  $F$  as the set

$$\{(x, f x). x \in F\}$$

So we are modeling partial functions by specifying the domain and the mapping function. We use the term “function” also for its graph.

```

type-synonym 'a graph = ('a  $\times$  real) set

```

```

definition graph :: 'a set  $\Rightarrow$  ('a  $\Rightarrow$  real)  $\Rightarrow$  'a graph
  where graph F f = {(x, f x) | x. x  $\in$  F}

```

```

lemma graphI [intro]: x  $\in$  F  $\Longrightarrow$  (x, f x)  $\in$  graph F f
  <proof>

```

```

lemma graphI2 [intro?]: x  $\in$  F  $\Longrightarrow$   $\exists$  t  $\in$  graph F f. t = (x, f x)
  <proof>

```

```

lemma graphE [elim?]:
  assumes (x, y)  $\in$  graph F f
  obtains x  $\in$  F and y = f x
  <proof>

```

### 7.2 Functions ordered by domain extension

A function  $h'$  is an extension of  $h$ , iff the graph of  $h$  is a subset of the graph of  $h'$ .

```

lemma graph-extI:
  ( $\bigwedge$ x. x  $\in$  H  $\Longrightarrow$  h x = h' x)  $\Longrightarrow$  H  $\subseteq$  H'
   $\Longrightarrow$  graph H h  $\subseteq$  graph H' h'
  <proof>

```

```

lemma graph-extD1 [dest?]: graph H h  $\subseteq$  graph H' h'  $\Longrightarrow$  x  $\in$  H  $\Longrightarrow$  h x = h' x
  <proof>

```

```

lemma graph-extD2 [dest?]: graph H h  $\subseteq$  graph H' h'  $\Longrightarrow$  H  $\subseteq$  H'
  <proof>

```

### 7.3 Domain and function of a graph

The inverse functions to *graph* are *domain* and *funct*.

**definition** *domain* :: 'a graph  $\Rightarrow$  'a set  
**where** *domain*  $g = \{x. \exists y. (x, y) \in g\}$

**definition** *funct* :: 'a graph  $\Rightarrow$  ('a  $\Rightarrow$  real)  
**where** *funct*  $g = (\lambda x. (SOME y. (x, y) \in g))$

The following lemma states that  $g$  is the graph of a function if the relation induced by  $g$  is unique.

**lemma** *graph-domain-funct*:  
**assumes** *uniq*:  $\bigwedge x y z. (x, y) \in g \Longrightarrow (x, z) \in g \Longrightarrow z = y$   
**shows** *graph* (*domain*  $g$ ) (*funct*  $g$ ) =  $g$   
 <proof>

### 7.4 Norm-preserving extensions of a function

Given a linear form  $f$  on the space  $F$  and a seminorm  $p$  on  $E$ . The set of all linear extensions of  $f$ , to superspaces  $H$  of  $F$ , which are bounded by  $p$ , is defined as follows.

**definition**  
*norm-pres-extensions* ::  
 'a::{plus, minus, uminus, zero} set  $\Rightarrow$  ('a  $\Rightarrow$  real)  $\Rightarrow$  'a set  $\Rightarrow$  ('a  $\Rightarrow$  real)  
 $\Rightarrow$  'a graph set

**where**  
*norm-pres-extensions*  $E p F f$   
 =  $\{g. \exists H h. g = \text{graph } H h$   
 $\wedge \text{linearform } H h$   
 $\wedge H \trianglelefteq E$   
 $\wedge F \trianglelefteq H$   
 $\wedge \text{graph } F f \subseteq \text{graph } H h$   
 $\wedge (\forall x \in H. h x \leq p x)\}$

**lemma** *norm-pres-extensionE* [elim]:  
**assumes**  $g \in \text{norm-pres-extensions } E p F f$   
**obtains**  $H h$   
**where**  $g = \text{graph } H h$   
**and** *linearform*  $H h$   
**and**  $H \trianglelefteq E$   
**and**  $F \trianglelefteq H$   
**and**  $\text{graph } F f \subseteq \text{graph } H h$   
**and**  $\forall x \in H. h x \leq p x$   
 <proof>

**lemma** *norm-pres-extensionI2* [intro]:  
*linearform*  $H h \Longrightarrow H \trianglelefteq E \Longrightarrow F \trianglelefteq H$   
 $\Longrightarrow \text{graph } F f \subseteq \text{graph } H h \Longrightarrow \forall x \in H. h x \leq p x$   
 $\Longrightarrow \text{graph } H h \in \text{norm-pres-extensions } E p F f$   
 <proof>

**lemma** *norm-pres-extensionI*:  
 $\exists H h. g = \text{graph } H h$

```

 $\wedge$  linearform  $H h$ 
 $\wedge H \leq E$ 
 $\wedge F \leq H$ 
 $\wedge \text{graph } F f \subseteq \text{graph } H h$ 
 $\wedge (\forall x \in H. h x \leq p x) \implies g \in \text{norm-pres-extensions } E p F f$ 
<proof>

```

end

## 8 The norm of a function

```

theory Function-Norm
imports Normed-Space Function-Order
begin

```

### 8.1 Continuous linear forms

A linear form  $f$  on a normed vector space  $(V, \|\cdot\|)$  is *continuous*, iff it is bounded, i.e.

$$\exists c \in \mathbb{R}. \forall x \in V. |f x| \leq c \cdot \|x\|$$

In our application no other functions than linear forms are considered, so we can define continuous linear forms as bounded linear forms:

```

locale continuous = var-V + norm-syntax + linearform +
  assumes bounded:  $\exists c. \forall x \in V. |f x| \leq c * \|x\|$ 

```

```

declare continuous.intro [intro?] continuous-axioms.intro [intro?]

```

```

lemma continuousI [intro]:
  fixes norm ::  $- \Rightarrow \text{real}$  ( $\|\cdot\|$ )
  assumes linearform  $V f$ 
  assumes  $r: \bigwedge x. x \in V \implies |f x| \leq c * \|x\|$ 
  shows continuous  $V \text{norm } f$ 
<proof>

```

### 8.2 The norm of a linear form

The least real number  $c$  for which holds

$$\forall x \in V. |f x| \leq c \cdot \|x\|$$

is called the *norm* of  $f$ .

For non-trivial vector spaces  $V \neq \{0\}$  the norm can be defined as

$$\|f\| = \sup_{x \neq 0} |f x| / \|x\|$$

For the case  $V = \{0\}$  the supremum would be taken from an empty set. Since  $\mathbb{R}$  is unbounded, there would be no supremum. To avoid this situation it must be guaranteed that there is an element in this set. This element must be  $\{ \} \geq 0$  so that *fn-norm* has the norm properties. Furthermore it does not have to

change the norm in all other cases, so it must be  $0$ , as all other elements are  $\{ \} \geq 0$ .

Thus we define the set  $B$  where the supremum is taken from as follows:

$$\{0\} \cup \{|f x| / \|x\|. \ x \neq 0 \wedge x \in F\}$$

$fn\text{-norm}$  is equal to the supremum of  $B$ , if the supremum exists (otherwise it is undefined).

```

locale fn-norm = norm-syntax +
  fixes  $B$  defines  $B \ V f \equiv \{0\} \cup \{|f x| / \|x\| \mid x. \ x \neq 0 \wedge x \in V\}$ 
  fixes  $fn\text{-norm}$  ( $\|\cdot\|$ --  $[0, 1000]$  999)
  defines  $\|f\|$ - $V \equiv \bigsqcup (B \ V f)$ 

```

```

locale normed-vectorspace-with-fn-norm = normed-vectorspace + fn-norm

```

```

lemma (in fn-norm) B-not-empty [intro]:  $0 \in B \ V f$ 
  <proof>

```

The following lemma states that every continuous linear form on a normed space  $(V, \|\cdot\|)$  has a function norm.

```

lemma (in normed-vectorspace-with-fn-norm) fn-norm-works:
  assumes continuous  $V$  norm  $f$ 
  shows lub ( $B \ V f$ ) ( $\|f\|$ - $V$ )
  <proof>

```

```

lemma (in normed-vectorspace-with-fn-norm) fn-norm-ub [iff?]:
  assumes continuous  $V$  norm  $f$ 
  assumes  $b$ :  $b \in B \ V f$ 
  shows  $b \leq \|f\|$ - $V$ 
  <proof>

```

```

lemma (in normed-vectorspace-with-fn-norm) fn-norm-leastB:
  assumes continuous  $V$  norm  $f$ 
  assumes  $b$ :  $\bigwedge b. \ b \in B \ V f \implies b \leq y$ 
  shows  $\|f\|$ - $V \leq y$ 
  <proof>

```

The norm of a continuous function is always  $\geq 0$ .

```

lemma (in normed-vectorspace-with-fn-norm) fn-norm-ge-zero [iff]:
  assumes continuous  $V$  norm  $f$ 
  shows  $0 \leq \|f\|$ - $V$ 
  <proof>

```

The fundamental property of function norms is:

$$|f x| \leq \|f\| \cdot \|x\|$$

```

lemma (in normed-vectorspace-with-fn-norm) fn-norm-le-cong:
  assumes continuous  $V$  norm  $f$  linearform  $V$   $f$ 
  assumes  $x$ :  $x \in V$ 
  shows  $|f x| \leq \|f\|$ - $V \ * \ \|x\|$ 
  <proof>

```

The function norm is the least positive real number for which the following inequation holds:

$$|f x| \leq c \cdot \|x\|$$

**lemma** (in *normed-vectorspace-with-fn-norm*) *fn-norm-least* [*intro?*]:

**assumes** *continuous V norm f*

**assumes** *ineq:  $\forall x \in V. |f x| \leq c * \|x\|$  and ge:  $0 \leq c$*

**shows**  $\|f\| - V \leq c$

*<proof>*

**end**

## 9 Zorn's Lemma

**theory** *Zorn-Lemma*

**imports** *~/src/HOL/Library/Zorn*

**begin**

Zorn's Lemmas states: if every linear ordered subset of an ordered set  $S$  has an upper bound in  $S$ , then there exists a maximal element in  $S$ . In our application,  $S$  is a set of sets ordered by set inclusion. Since the union of a chain of sets is an upper bound for all elements of the chain, the conditions of Zorn's lemma can be modified: if  $S$  is non-empty, it suffices to show that for every non-empty chain  $c$  in  $S$  the union of  $c$  also lies in  $S$ .

**theorem** *Zorn's-Lemma*:

**assumes**  $r: \bigwedge c. c \in \text{chain } S \implies \exists x. x \in c \implies \bigcup c \in S$

**and**  $aS: a \in S$

**shows**  $\exists y \in S. \forall z \in S. y \subseteq z \implies y = z$

*<proof>*

**end**

## Part II

# Lemmas for the Proof

## 10 The supremum w.r.t. the function order

**theory** *Hahn-Banach-Sup-Lemmas*  
**imports** *Function-Norm Zorn-Lemma*  
**begin**

This section contains some lemmas that will be used in the proof of the Hahn-Banach Theorem. In this section the following context is presumed. Let  $E$  be a real vector space with a seminorm  $p$  on  $E$ .  $F$  is a subspace of  $E$  and  $f$  a linear form on  $F$ . We consider a chain  $c$  of norm-preserving extensions of  $f$ , such that  $\bigcup c = \text{graph } H h$ . We will show some properties about the limit function  $h$ , i.e. the supremum of the chain  $c$ .

Let  $c$  be a chain of norm-preserving extensions of the function  $f$  and let  $\text{graph } H h$  be the supremum of  $c$ . Every element in  $H$  is member of one of the elements of the chain.

**lemmas**  $[\text{dest?}] = \text{chainD}$   
**lemmas**  $\text{chainE2} [\text{elim?}] = \text{chainD2} [\text{elim-format, standard}]$

**lemma** *some- $H'h'$* :

**assumes**  $M: M = \text{norm-pres-extensions } E p F f$   
**and**  $cM: c \in \text{chain } M$   
**and**  $u: \text{graph } H h = \bigcup c$   
**and**  $x: x \in H$   
**shows**  $\exists H' h'. \text{graph } H' h' \in c$   
 $\wedge (x, h x) \in \text{graph } H' h'$   
 $\wedge \text{linearform } H' h' \wedge H' \trianglelefteq E$   
 $\wedge F \trianglelefteq H' \wedge \text{graph } F f \subseteq \text{graph } H' h'$   
 $\wedge (\forall x \in H'. h' x \leq p x)$

*<proof>*

Let  $c$  be a chain of norm-preserving extensions of the function  $f$  and let  $\text{graph } H h$  be the supremum of  $c$ . Every element in the domain  $H$  of the supremum function is member of the domain  $H'$  of some function  $h'$ , such that  $h$  extends  $h'$ .

**lemma** *some- $H'h'$* :

**assumes**  $M: M = \text{norm-pres-extensions } E p F f$   
**and**  $cM: c \in \text{chain } M$   
**and**  $u: \text{graph } H h = \bigcup c$   
**and**  $x: x \in H$   
**shows**  $\exists H' h'. x \in H' \wedge \text{graph } H' h' \subseteq \text{graph } H h$   
 $\wedge \text{linearform } H' h' \wedge H' \trianglelefteq E \wedge F \trianglelefteq H'$   
 $\wedge \text{graph } F f \subseteq \text{graph } H' h' \wedge (\forall x \in H'. h' x \leq p x)$

*<proof>*

Any two elements  $x$  and  $y$  in the domain  $H$  of the supremum function  $h$  are both in the domain  $H'$  of some function  $h'$ , such that  $h$  extends  $h'$ .

**lemma** *some- $H'h'2$* :

**assumes**  $M$ :  $M = \text{norm-pres-extensions } E \ p \ F \ f$   
**and**  $cM$ :  $c \in \text{chain } M$   
**and**  $u$ :  $\text{graph } H \ h = \bigcup c$   
**and**  $x$ :  $x \in H$   
**and**  $y$ :  $y \in H$   
**shows**  $\exists H' \ h', x \in H' \wedge y \in H'$   
 $\wedge \text{graph } H' \ h' \subseteq \text{graph } H \ h$   
 $\wedge \text{linearform } H' \ h' \wedge H' \leq E \wedge F \leq H'$   
 $\wedge \text{graph } F \ f \subseteq \text{graph } H' \ h' \wedge (\forall x \in H'. h' \ x \leq p \ x)$

*<proof>*

The relation induced by the graph of the supremum of a chain  $c$  is definite, i. e.  $t$  is the graph of a function.

**lemma** *sup-definite*:

**assumes**  $M\text{-def}$ :  $M \equiv \text{norm-pres-extensions } E \ p \ F \ f$   
**and**  $cM$ :  $c \in \text{chain } M$   
**and**  $xy$ :  $(x, y) \in \bigcup c$   
**and**  $xz$ :  $(x, z) \in \bigcup c$

**shows**  $z = y$

*<proof>*

The limit function  $h$  is linear. Every element  $x$  in the domain of  $h$  is in the domain of a function  $h'$  in the chain of norm preserving extensions. Furthermore,  $h$  is an extension of  $h'$  so the function values of  $x$  are identical for  $h'$  and  $h$ . Finally, the function  $h'$  is linear by construction of  $M$ .

**lemma** *sup-lf*:

**assumes**  $M$ :  $M = \text{norm-pres-extensions } E \ p \ F \ f$   
**and**  $cM$ :  $c \in \text{chain } M$   
**and**  $u$ :  $\text{graph } H \ h = \bigcup c$   
**shows**  $\text{linearform } H \ h$

*<proof>*

The limit of a non-empty chain of norm preserving extensions of  $f$  is an extension of  $f$ , since every element of the chain is an extension of  $f$  and the supremum is an extension for every element of the chain.

**lemma** *sup-ext*:

**assumes**  $\text{graph}$ :  $\text{graph } H \ h = \bigcup c$   
**and**  $M$ :  $M = \text{norm-pres-extensions } E \ p \ F \ f$   
**and**  $cM$ :  $c \in \text{chain } M$   
**and**  $ex$ :  $\exists x. x \in c$   
**shows**  $\text{graph } F \ f \subseteq \text{graph } H \ h$

*<proof>*

The domain  $H$  of the limit function is a superspace of  $F$ , since  $F$  is a subset of  $H$ . The existence of the  $0$  element in  $F$  and the closure properties follow from the fact that  $F$  is a vector space.

**lemma** *sup-supF*:

**assumes**  $\text{graph}$ :  $\text{graph } H \ h = \bigcup c$   
**and**  $M$ :  $M = \text{norm-pres-extensions } E \ p \ F \ f$

```

and  $cM: c \in \text{chain } M$ 
and  $ex: \exists x. x \in c$ 
and  $FE: F \leq E$ 
shows  $F \leq H$ 
<proof>

```

The domain  $H$  of the limit function is a subspace of  $E$ .

```

lemma sup-subE:
assumes  $graph: \text{graph } H \ h = \bigcup c$ 
and  $M: M = \text{norm-pres-extensions } E \ p \ F \ f$ 
and  $cM: c \in \text{chain } M$ 
and  $ex: \exists x. x \in c$ 
and  $FE: F \leq E$ 
and  $E: \text{vectorspace } E$ 
shows  $H \leq E$ 
<proof>

```

The limit function is bounded by the norm  $p$  as well, since all elements in the chain are bounded by  $p$ .

```

lemma sup-norm-pres:
assumes  $graph: \text{graph } H \ h = \bigcup c$ 
and  $M: M = \text{norm-pres-extensions } E \ p \ F \ f$ 
and  $cM: c \in \text{chain } M$ 
shows  $\forall x \in H. h \ x \leq p \ x$ 
<proof>

```

The following lemma is a property of linear forms on real vector spaces. It will be used for the lemma *abs-Hahn-Banach* (see page ??). For real vector spaces the following inequations are equivalent:

$$\forall x \in H. |h \ x| \leq p \ x \quad \text{and} \quad \forall x \in H. h \ x \leq p \ x$$

```

lemma abs-ineq-iff:
assumes  $\text{subspace } H \ E$  and  $\text{vectorspace } E$  and  $\text{seminorm } E \ p$ 
and  $\text{linearform } H \ h$ 
shows  $(\forall x \in H. |h \ x| \leq p \ x) = (\forall x \in H. h \ x \leq p \ x)$  (is ?L = ?R)
<proof>

```

**end**

## 11 Extending non-maximal functions

```

theory Hahn-Banach-Ext-Lemmas
imports Function-Norm
begin

```

In this section the following context is presumed. Let  $E$  be a real vector space with a seminorm  $q$  on  $E$ .  $F$  is a subspace of  $E$  and  $f$  a linear function on  $F$ . We consider a subspace  $H$  of  $E$  that is a superspace of  $F$  and a linear form  $h$  on  $H$ .  $H$  is not equal to  $E$  and  $x_0$  is an element in  $E - H$ .  $H$  is extended to the

direct sum  $H' = H + \text{lin } x_0$ , so for any  $x \in H'$  the decomposition of  $x = y + a \cdot x$  with  $y \in H$  is unique.  $h'$  is defined on  $H'$  by  $h' x = h y + a \cdot \xi$  for a certain  $\xi$ .

Subsequently we show some properties of this extension  $h'$  of  $h$ .

This lemma will be used to show the existence of a linear extension of  $f$  (see page ??). It is a consequence of the completeness of  $\mathbb{R}$ . To show

$$\exists \xi. \forall y \in F. a y \leq \xi \wedge \xi \leq b y$$

it suffices to show that

$$\forall u \in F. \forall v \in F. a u \leq b v$$

**lemma** *ex-xi*:

**assumes** *vectorspace*  $F$

**assumes**  $r: \bigwedge u v. u \in F \implies v \in F \implies a u \leq b v$

**shows**  $\exists xi::real. \forall y \in F. a y \leq xi \wedge xi \leq b y$

*<proof>*

The function  $h'$  is defined as a  $h' x = h y + a \cdot \xi$  where  $x = y + a \cdot \xi$  is a linear extension of  $h$  to  $H'$ .

**lemma** *h'-lf*:

**assumes** *h'-def*:  $h' \equiv \lambda x. \text{let } (y, a) =$

*SOME*  $(y, a). x = y + a \cdot x_0 \wedge y \in H \text{ in } h y + a * xi$

**and** *H'-def*:  $H' \equiv H \oplus \text{lin } x_0$

**and** *HE*:  $H \trianglelefteq E$

**assumes** *linearform*  $H h$

**assumes** *x0*:  $x_0 \notin H \ x_0 \in E \ x_0 \neq 0$

**assumes** *E*: *vectorspace*  $E$

**shows** *linearform*  $H' h'$

*<proof>*

The linear extension  $h'$  of  $h$  is bounded by the seminorm  $p$ .

**lemma** *h'-norm-pres*:

**assumes** *h'-def*:  $h' \equiv \lambda x. \text{let } (y, a) =$

*SOME*  $(y, a). x = y + a \cdot x_0 \wedge y \in H \text{ in } h y + a * xi$

**and** *H'-def*:  $H' \equiv H \oplus \text{lin } x_0$

**and** *x0*:  $x_0 \notin H \ x_0 \in E \ x_0 \neq 0$

**assumes** *E*: *vectorspace*  $E$  **and** *HE*: *subspace*  $H E$

**and** *seminorm*  $E p$  **and** *linearform*  $H h$

**assumes** *a*:  $\forall y \in H. h y \leq p y$

**and** *a'*:  $\forall y \in H. -p (y + x_0) - h y \leq xi \wedge xi \leq p (y + x_0) - h y$

**shows**  $\forall x \in H'. h' x \leq p x$

*<proof>*

**end**

## Part III

# The Main Proof

## 12 The Hahn-Banach Theorem

**theory** *Hahn-Banach*  
**imports** *Hahn-Banach-Lemmas*  
**begin**

We present the proof of two different versions of the Hahn-Banach Theorem, closely following [1, §36].

### 12.1 The Hahn-Banach Theorem for vector spaces

**Hahn-Banach Theorem.** Let  $F$  be a subspace of a real vector space  $E$ , let  $p$  be a semi-norm on  $E$ , and  $f$  be a linear form defined on  $F$  such that  $f$  is bounded by  $p$ , i.e.  $\forall x \in F. f x \leq p x$ . Then  $f$  can be extended to a linear form  $h$  on  $E$  such that  $h$  is norm-preserving, i.e.  $h$  is also bounded by  $p$ .

#### Proof Sketch.

1. Define  $M$  as the set of norm-preserving extensions of  $f$  to subspaces of  $E$ . The linear forms in  $M$  are ordered by domain extension.
2. We show that every non-empty chain in  $M$  has an upper bound in  $M$ .
3. With Zorn's Lemma we conclude that there is a maximal function  $g$  in  $M$ .
4. The domain  $H$  of  $g$  is the whole space  $E$ , as shown by classical contradiction:
  - Assuming  $g$  is not defined on whole  $E$ , it can still be extended in a norm-preserving way to a super-space  $H'$  of  $H$ .
  - Thus  $g$  can not be maximal. Contradiction!

**theorem** *Hahn-Banach*:

**assumes**  $E$ : *vectorspace*  $E$  **and** *subspace*  $F$   $E$

**and** *seminorm*  $E$   $p$  **and** *linearform*  $F$   $f$

**assumes**  $fp$ :  $\forall x \in F. f x \leq p x$

**shows**  $\exists h. linearform$   $E$   $h \wedge (\forall x \in F. h x = f x) \wedge (\forall x \in E. h x \leq p x)$

— Let  $E$  be a vector space,  $F$  a subspace of  $E$ ,  $p$  a seminorm on  $E$ ,

— and  $f$  a linear form on  $F$  such that  $f$  is bounded by  $p$ ,

— then  $f$  can be extended to a linear form  $h$  on  $E$  in a norm-preserving way.

*<proof>*

### 12.2 Alternative formulation

The following alternative formulation of the Hahn-Banach Theorem uses the fact that for a real linear form  $f$  and a seminorm  $p$  the following inequations are equivalent:<sup>1</sup>

<sup>1</sup>This was shown in lemma *abs-ineq-iff* (see page 21).

$$\forall x \in H. |h x| \leq p x \quad \text{and} \quad \forall x \in H. h x \leq p x$$

**theorem** *abs-Hahn-Banach*:

**assumes**  $E$ : *vectorspace*  $E$  **and**  $FE$ : *subspace*  $F E$

**and**  $lf$ : *linearform*  $F f$  **and**  $sn$ : *seminorm*  $E p$

**assumes**  $fp$ :  $\forall x \in F. |f x| \leq p x$

**shows**  $\exists g$ . *linearform*  $E g$

$\wedge (\forall x \in F. g x = f x)$

$\wedge (\forall x \in E. |g x| \leq p x)$

*<proof>*

### 12.3 The Hahn-Banach Theorem for normed spaces

Every continuous linear form  $f$  on a subspace  $F$  of a norm space  $E$ , can be extended to a continuous linear form  $g$  on  $E$  such that  $\|f\| = \|g\|$ .

**theorem** *norm-Hahn-Banach*:

**fixes**  $V$  **and** *norm*  $(\|-|\|)$

**fixes**  $B$  **defines**  $\bigwedge V f. B V f \equiv \{0\} \cup \{|f x| / \|x\| \mid x. x \neq 0 \wedge x \in V\}$

**fixes**  $fn$ -*norm*  $(\|-|\|$ -  $[0, 1000]$  999)

**defines**  $\bigwedge V f. \|f\|$ - $V \equiv \bigsqcup (B V f)$

**assumes**  $E$ -*norm*: *normed-vectorspace*  $E$  *norm* **and**  $FE$ : *subspace*  $F E$

**and** *linearform*: *linearform*  $F f$  **and** *continuous*  $F$  *norm*  $f$

**shows**  $\exists g$ . *linearform*  $E g$

$\wedge$  *continuous*  $E$  *norm*  $g$

$\wedge (\forall x \in F. g x = f x)$

$\wedge \|g\|$ - $E = \|f\|$ - $F$

*<proof>*

**end**

## References

- [1] H. Heuser. *Funktionalanalysis: Theorie und Anwendung*. Teubner, 1986.
- [2] L. Narici and E. Beckenstein. The Hahn-Banach Theorem: The life and times. In *Topology Atlas*. York University, Toronto, Ontario, Canada, 1996. <http://at.yorku.ca/topology/preprint.htm> and <http://at.yorku.ca/p/a/a/a/16.htm>.
- [3] B. Nowak and A. Trybulec. Hahn-Banach theorem. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.uwb.edu.pl/JFM/Vol5/hahnban.html>.