

Chapter 1

Maps

File: foundations/sections/06_maps.ftl.tex

[readtex foundations/sections/04_pairs-and-products.ftl.tex]

1.1 Ranges

FOUNDATIONS_06_4284980337311744

Definition 1.1. Let f be a map. A value of f is an object b such that $b = f(a)$ for some $a \in \text{dom}(f)$.

FOUNDATIONS_06_1938831225913344

Definition 1.2. Let f be a map. The range of f is

$$\{f(a) \mid a \in \text{dom}(f)\}.$$

Let $\text{range}(f)$ stand for the range of f .

FOUNDATIONS_06_6386349418479616

Proposition 1.3. Let f be a map and b be an object. b is a value of f iff $b \in \text{range}(f)$.

Proof. Case b is a value of f . Take $a \in \text{dom}(f)$ such that $b = f(a)$. b is an object. Hence $b \in \text{range}(f)$. End.

Case $b \in \text{range}(f)$. Then b is an object such that $b = f(a)$ for some $a \in \text{dom}(f)$. Hence b is a value of f . End. \square

1.2 The identity map

FOUNDATIONS_06_1920902360989696

Definition 1.4. Let A be a class. id_A is the map h such that h is defined on A and $h(a) = a$ for all $a \in A$.

Let the identity map on A stand for id_A .

1.3 Composition

FOUNDATIONS_06_7605717729017856

Definition 1.5. Let f, g be maps. Assume $\text{range}(f) \subseteq \text{dom}(g)$. $g \circ f$ is the map h such that h is defined on $\text{dom}(f)$ and $h(a) = g(f(a))$ for all $a \in \text{dom}(f)$.

Let the composition of g and f stand for $g \circ f$.

1.4 Restriction

FOUNDATIONS_06_7095412741636096

Definition 1.6. Let f be a map and $X \subseteq \text{dom}(f)$. $f \upharpoonright X$ is the map h such that h is defined on X and $h(a) = f(a)$ for all $a \in X$.

Let the restriction of f to X stand for $f \upharpoonright X$.

FOUNDATIONS_06_2170189258948608

Proposition 1.7. Let A be a class and $X \subseteq A$. Then $\text{id}_A \upharpoonright X = \text{id}_X$.

1.5 Images and preimages

FOUNDATIONS_06_3038237683613696

Definition 1.8. Let f be a map and A be a class. The image of A under f is

$$\{f(a) \mid a \in \text{dom}(f) \cap A\}.$$

Let the direct image of A under f stand for the image of A under f . Let $f_*(A)$ stand for the image of A under f .

Let $f[A]$ stand for $f_*(A)$.

FOUNDATIONS_06_4563167805964288

Definition 1.9. Let f be a map and B be a class. The preimage of B under f is

$$\{a \in \text{dom}(f) \mid f(a) \in B\}.$$

Let the inverse image of B under f stand for the preimage of B under f . Let $f^*(B)$ stand for the preimage of B under f .

1.6 Maps between classes

FOUNDATIONS_06_6934038600220672

Definition 1.10. Let A be a class. A map of A is a map f such that $\text{dom}(f) = A$.

FOUNDATIONS_06_7725375157174272

Definition 1.11. Let B be a class. A map to B is a map f such that $f(a) \in B$ for each $a \in \text{dom}(f)$.

FOUNDATIONS_06_2823507398361088

Definition 1.12. Let A, B be classes. A map from A to B is a map f such that $\text{dom}(f) = A$ and $f(a) \in B$ for each $a \in A$.

Let $f : A \rightarrow B$ stand for f is a map from A to B .

FOUNDATIONS_06_3390734908522496

Definition 1.13. Let A be a class. A map on A is a map from A to A .

FOUNDATIONS_06_3312973569327104

Proposition 1.14. Let A, B be classes and $f, g : A \rightarrow B$. Assume that $f(a) = g(a)$ for all $a \in A$. Then $f = g$.

Proposition 1.15. Let A, B be classes and f be a map of A . Assume that $f(a) \in B$ for all $a \in A$. Then f is a map from A to B iff $\text{range}(f) \subseteq B$.

FOUNDATIONS_06_5104361690628096

Proposition 1.16. Let A be a class. Then id_A is a map on A .

FOUNDATIONS_06_1706446651654144

Proposition 1.17. Let A, B, C be classes and $f : A \rightarrow B$ and $g : B \rightarrow C$. Then $g \circ f : A \rightarrow C$.

FOUNDATIONS_06_4078561256275968

Proposition 1.18. Let A, B be classes and $f : A \rightarrow B$ and $X \subseteq A$. Then $f \upharpoonright X : X \rightarrow B$.

FOUNDATIONS_06_3964401904254976

Proposition 1.19. Let A, B be classes and $f : A \rightarrow B$. Then

$$f \circ \text{id}_A = f = \text{id}_B \circ f.$$

Proof. A is the domain of $f \circ \text{id}_A$ and the domain of f and the domain of $\text{id}_B \circ f$. We have $(f \circ \text{id}_A)(a) = f(\text{id}_A(a)) = f(a) = \text{id}_B(f(a)) = (\text{id}_B \circ f)(a)$ for all $a \in A$. Hence

$$f \circ \text{id}_A = f = \text{id}_B \circ f.$$

□

FOUNDATIONS_06_3118771061391360

Proposition 1.20. Let A be a class and $X \subseteq A$. Then

$$\text{id}_A \upharpoonright X = \text{id}_X.$$

Proof. We have $\text{dom}(\text{id}_A \upharpoonright X) = X = \text{dom}(\text{id}_X)$. $(\text{id}_A \upharpoonright X)(a) = \text{id}_A(a) = a = \text{id}_X(a)$ for all $a \in X$. Hence $\text{id}_A \upharpoonright X = \text{id}_X$. □

FOUNDATIONS_06_6866147389472768

Proposition 1.21. Let A, B, C, D be classes and $f : A \rightarrow B$ and $g : B \rightarrow C$ and $h : C \rightarrow D$. Then

$$h \circ (g \circ f) = (h \circ g) \circ f.$$

Proof. $h \circ (g \circ f)$ and $(h \circ g) \circ f$ are maps from A to D .

Let us show that $(h \circ (g \circ f))(a) = ((h \circ g) \circ f)(a)$ for all $a \in A$. Let $a \in A$. Then $(h \circ (g \circ f))(a) = h((g \circ f)(a)) = h(g(f(a))) = (h \circ g)(f(a)) = ((h \circ g) \circ f)(a)$. End.

Hence $h \circ (g \circ f) = (h \circ g) \circ f$. □

1.7 Maps and products

FOUNDATIONS_06_5135634870042624

Definition 1.22. Let f be a map such that $\text{dom}(f) = A \times B$ for some nonempty classes A, B . Let a be an object such that $(a, b) \in \text{dom}(f)$ for some object b . $f(a, -)$ is the map such that $\text{dom}(f(a, -)) = B$ and $f(a, -)(b) = f(a, b)$ for all $b \in B$ where B is the class such that $\text{dom}(f) = A \times B$ for some class A .

FOUNDATIONS_06_3621991366000640

Definition 1.23. Let f be a map such that $\text{dom}(f) = A \times B$ for some nonempty classes A, B . Let b be an object such that $(a, b) \in \text{dom}(f)$ for some object a . $f(-, b)$ is the map such that $\text{dom}(f(-, b)) = A$ and $f(-, b)(a) = f(a, b)$ for all $a \in A$ where A is the class such that $\text{dom}(f) = A \times B$ for some class B .

FOUNDATIONS_06_8946256734846976

Proposition 1.24. Let A, B, C be classes such that A, B are nonempty and $a \in A$. Let f be a map from $A \times B$ to C . Then $f(a, -)$ is a map from B to C .

FOUNDATIONS_06_8080207992848384

Proposition 1.25. Let A, B, C be classes such that A, B are nonempty and $b \in B$. Let f be a map from $A \times B$ to C . Then $f(-, b)$ is a map from A to C .

FOUNDATIONS_06_2754759509409792

Proposition 1.26. Let A, B, C be classes and f be a map of $A \times B$. Assume that $f(a, b) \in C$ for all $a \in A$ and all $b \in B$. Then f is a map from $A \times B$ to C .

FOUNDATIONS_06_2304295212941312

Proposition 1.27. Let A, B, C be classes and f be a map from $A \times B$ to C . Let $a \in A$ and $b \in B$. Then $f(a, b) \in C$.